SPRINGBROOK BIO-PHYSICAL

BIO-PHYSICAL CAPABILITIES FOR WILDLIFE

— MOOSE —

DESCRIPTIVE LEGEND

On this reconnaissance map, the land is divided into units on a biophysical basis which takes into account geology, landform, relief, soils and vegetation The biophysical framework is the same as that used for capability inventories of forestry, agriculture, native forage and recreation in the map area Base data maps of soils, landforms, vegetation and climate as well as biophysical capability maps for forestry, recreation and agriculture are available from the Resource Analysis Unit, E L U C Secretariat, Victoria

CAPABILITY CLASSIFICATIONS

Capability ratings are established on the ability of the land to meet the total needs of the wildlife species over the long term In terms of food and cover needs, the ratings are based on the optimum vegetational stage (successional stage) that can be maintained assuming good management practices that do not degrade the environment or land base The application of management prescriptions are limited to the following non-intensive practices:

1) Prescribed burning or grazing
2) Prescribed logging or occasional slashing
3) Protection from fire or any land-use practice that would reduce the productivity of the land for the wildlife species

When assigning capability ratings, map units are evaluated on the basis of their position in the ecosystem as a whole and not in isolation

Capability ratings reflect only biological and physical parameters of the environment and do not take into account human elements such as social or economic parameters
Capability ratings assigned do not reflect present land use, ownership, lack of access, distance from cities, or amount of hunting pressure

Carrying capacity estimates based on the number of animals which can be supported annually on a sustained basis are assigned to each capability class as a measure of productivity

CLASS 1 LANDS IN THIS CLASS HAVE THE LEAST LIMITATION TO THE PRODUCTION OF MOOSE

They can provide a wide variety and abundance of food plants and other necessary habitat elements These very high capability lands can support an estimated 11 to 23 moose per square mile annually

When required, this class may be subdivided on the basis of productivity into Class 1a (11 to 15 moose per square mile per year), Class 1b (15 to 19 moose per square mile per year) or Class 1c (19 to 23 moose per square mile per year) etc using an increment equivalent to 25% of the upper figure for Class 1a

CLASS 2 LANDS IN THIS CLASS HAVE SLIGHT LIMITATIONS TO THE PRODUCTION OF MOOSE

Capability on these lands is high, but less than Class 1 These lands can support an estimated 8 to 11 moose per square mile annually

CLASS 3 LANDS IN THIS CLASS HAVE MODERATE LIMITATIONS TO THE PRODUCTION OF MOOSE

Capability on these lands is moderate to moderately high and they can support an estimated 4 to 8 moose per square mile annually

CLASS 4 LANDS IN THIS CLASS HAVE MODERATELY SEVERE LIMITATIONS TO THE PRODUCTION OF MOOSE

Capability on these lands is moderate to moderately low and they can support an estimated 1 to 4 moose per square mile annually

CLASS 5 LANDS IN THIS CLASS HAVE SEVERE LIMITATIONS TO THE PRODUCTION OF MOOSE

Capability on these lands is very low and it is estimated they can support less than 1 moose per square mile annually

CLASS 6 LANDS IN THIS CLASS HAVE LIMITATIONS SO SEVERE THAT THERE IS NO PRODUCTION OF MOOSE

L C low capability - this designation applies to extensive areas which are low or lacking in habitat potential for a species and are therefore excluded from classification It is also applied to lands on which man's activities have more or less permanently obscured or eliminated the habitat potential for a species

SUBCLASSES

The most significant environmental factors limiting the production of the species and thus determining the capability class, are indicated on the map by subclasses In most cases, the limitations do not affect the animals themselves, but rather the ability of the land to meet the needs of the species in terms of food, cover, and other requirements For convenience, the subclasses are placed in three main categories those relating to climate, those relating to the inherent characteristics of the land and those relating to the vegetative cover which, although reflecting characteristics of the land and/or climate, are not adequately covered or explained by limiting subclasses in these categories

LIMITATIONS OF CLIMATE

SUBCLASS A aridity - Drought or aridity that adversely affects the habitat

SUBCLASS C climate - A combination of climatic factors acting to reduce favourable habitat and the production and survival of the ungulate species

SUBCLASS Q snow depth - Excessive snow depth that reduces the mobility of the ungulate species or the availability of food plants

SUBCLASS U exposure - Exposure to extreme climatic factors such as extreme or drying winds that adversely affect the species or its habitat

LIMITATIONS OF THE LAND

SUBCLASS E erodability - Maintenance of optimum vegetation for food and cover restricted at assumed management level due to susceptability to soil erosion ranging from sheet erosion through minor gullying

SUBCLASS F fertility - Lack of available nutrients in the soil for optimum plant growth

SUBCLASS G landform Poor distribution of interspersion of landforms necessary for optimum habitat

SUBCLASS I inundation - Excessive water level fluctuation or tidal action that adversely affects the habitat or survival of the species

SUBCLASS M soil moisture - Poor soil moisture, either excessive of deficient

SUBCLASS N soil chemistry - Excessive salinity, lack of essential trace elements or abundance of toxic elements in the soil

SUBCLASS R soil depth - Restriction of the rooting zone by bedrock or other impervious layers

SUBCLASS S soil sensitivity - Maintenance of optimum vegetation restricted at assumed management level due to susceptibility to loss of soil nutrients through leaching

SUBCLASS T adverse topography - Either steepness or flatness of the land

SUBCLASS W mass wasting - Extensive erosion processes, active or potential, as in talus slope or slope failure situations

LIMITATIONS OF VEGETATION

SUBCLASS D dense regeneration - Maintenance of optimum vegetation for food and cover restricted a assumed management level by susceptibility to dense regeneration of vegetation

SUBCLASS V rapid seral succession - Maintenance of optimum vegetation for food and cover restricted at assumed management level by rapid seral succession Maintenance of a more favourable condition is either beyond assumed management levels or would result in impairment of the land base over time at assumed management levels

SPECIAL CLASS INDICATORS

The following are used to denote known habitat elements considered critical to the distribution or abundance of the species

Seasonal concentration areas are shown by the following symbols

W areas of winter concentrations
Sp areas of spring concentration
S areas of summer concentration
F areas of fall concentration
L denotes mineral licks used by the species

denotes an important movement corridor for the species

CONVENTIONS

Large arabic numerals denote capability class

Small lower case letters placed after capability class 1 denote a subdivision of that class

Large upper case letters placed after the capability class indicate the special class indicator

Small arabic numerals placed after class numeral, class subdivision letter or special class symbol indicate the approximate proportion (in tenths) of the complex that is represented by that class The dominant class appears first in the symbol

Small upper case letters placed after class numeral, class subdivision letter or special class indicator denotes the subclass

For a more detailed description of the classification system the reader should refer to the manual of Biophysical Capability Classsification for Wildlife (Ungulates) in British Columbia This manual presently at print will be available from the Resource Analysis Unit, E L U C Secretariat, Victoria, B C

EXAMPLES

An area of class 5 land with some depth, dense regeneration and landform limitations is shown thus

$5^{\frac{G}{D}}$

A class 1a fall concentration area with limitations due to snow depth is shown thus

$1aF^{Q}$

An area of which 70% is class 5 with limitations due to exposed bedrock and snow depth and 30% is class 3 wintering area with limitations due to snow depths and exposure to extreme winds is shown thus

$5\frac{7}{G}\text{-}3W\frac{3}{Q}$

GEOGRAPHIC DIVISION
ENVIRONMENT AND LAND USE COMMITTEE
SECRETARIAT
VICTORIA, B C

COMPILED BY A.J.L.	AMMENDMENTS
DRAWN BY G.L.E.	
DATE March 27, 1975	
CHECKED BY A.J.L.	PROJECT SPRINGBROOK
DATE April 14, 1975	SHEET 13 OF 27

307282