Sharp-tailed Snake

Endangered in Canada and Red-listed in British Columbia

The **Sharp-tailed Snake** is a small rare snake found on Vancouver Island and the Gulf Islands, British Columbia. This guide provides information on how to identify the snake and distinguish it from the more common garter snakes. All snakes found in this area are non-venomous and harmless to humans.

Sharp-tailed Snake (Contia tenuis)

SIZE Small and slender, worm-like; length of adults about 20–30 cm; length of young at hatching about 10 cm.

COLOUR Reddish or greyish brown; juveniles often brighter red than adults, similar to earthworms.

DISTINGUISHING FEATURES Sharp, thorn-like tail tip; black and white barring on the underside; smooth body scales; divided anal scale (see reverse side for details).

ACTIVITY Most active in late February to June and again in late September to October.

HABITS Usually hides under rocks, decaying wood, or debris; seldom seen out in the open.



BRITISH COLUMBIA Environment



Garter Snakes (Thamnophis species)

SPECIES IN BC Common Garter Snake (*T. sirtalis*); Western Terrestrial Garter Snake (*T. elegans*); Northwestern Garter Snake (*T. ordinoides*).

SIZE Length of adults up to about 66 cm (*T. ordinoides*), 100 cm (*T. elegans*), and 130 cm (*T. sirtalis*); young at birth about 15–20 cm long.

COLOUR Variable; often with red, yellow or orange stripes running along the sides and the mid-back on grey, brown or black background; some individuals lack stripes.

DISTINGUISHING FEATURES Body often striped; long tapering tail; keeled body scales resulting in rough texture.

ACTIVITY Most active in the spring and early summer but can be encountered any time, apart from cold periods in winter.

HABITS Often seen out in the open either sunning itself or moving through the undergrowth; shelters under rocks, decaying wood, and debris.



Northwestern Garter Snake

Common Garter Snake

Identification Key

(only applicable for southwestern BC)

